

# Hodge theory and Moduli

Phillip Griffiths\*

---

\*Clay Lecture, based in part on joint work with Mark Green, Radu Laza and Colleen Robles. Some the the lecture draws on the work of and discussions with Marco Franciosi, Rita Pardini and Sönke Rollenske. Some general references are [GGLR], [GG], [G] and [FPR]. Further specific references will be given in the lecture.

## Outline

- I. Introduction
  - II. Backgroup and general results
    - A. Moduli theory
    - B. Hodge theory
    - C. Some general results
  - III.  $I$ -surfaces
- References

# I. Introduction

- ▶ Moduli is a topic of central interest in algebraic geometry. The theory roughly organizes into three areas:
  - ▶ varieties  $X$  of general type ( $\kappa(X) = \dim X, K_X > 0$ );
  - ▶ Calabi-Yau varieties ( $\kappa(X) = 0, K_X = 0$ );
  - ▶ Fano varieties ( $\kappa(X) = -\infty, K_X < 0$ ).

This lecture is mainly concerned with the first type.

- ▶ The techniques for studying moduli also roughly divide into three types:
  - ▶ algebraic (birational geometry, singularity theory, geometric invariant theory (GIT), etc.);
  - ▶ Hodge theoretic (topological and geometric);
  - ▶ analytic ( $L^2$ - $\bar{\partial}$  techniques, construction and properties of special metrics).

The algebraic techniques are currently the dominant ones. The three methods also of course interact; e.g., complex analysis plays a central role in Hodge theory.

- ▶ For varieties of general type, drawing on ideas from the minimal model program Kollár–Shepherd-Baron–Alexeev (KSBA) proved the existence of a moduli space  $\mathcal{M}$  having a canonical completion, later proved to be projective (cf. [KSB], the survey paper [K] and the references cited therein).
- ▶ For this talk the general motivating question is

*What is the structure of  $\overline{\mathcal{M}}$ ?*

By structure, informally stated we mean the stratification of  $\overline{\mathcal{M}}$  where the strata correspond to varieties of the same deformation type (equisingular deformations). We also include the incidence relations among the strata. For surfaces where there is a classification, we include which surfaces in the classification occur in a stratum.

The method we will discuss to study this question is to use Hodge theory. With notation to be more fully explained later, we denote by  $\mathcal{P}$  the image of the period mapping  $\Phi : \mathcal{M} \rightarrow \Gamma \backslash D$  and by  $\overline{\mathcal{P}}$  the canonical completion of  $\mathcal{P}$ .<sup>†</sup> To help understand the structure there are two basic types of subvarieties of  $\overline{\mathcal{P}}$  and then there is the amalgam of these. The first type is the stratification associated to the boundary components given by limiting mixed Hodge structures  $(V, Q, W(N), F_{\text{lim}})$ <sup>‡</sup> that occur when polarized Hodge structures degenerate.

---

<sup>†</sup>Notably, in the non-classical case  $\overline{\mathcal{P}}$  is not a subset of a completion of  $\Gamma \backslash D$ ; it is a relative construction associated to period mapping.


<sup>‡</sup>The notations and terminology will be recalled below. For general background in Hodge theory we refer to Colleen Robles' lecture or to [GGLR] and [GG].

Roughly speaking one thinks of going in  $\mathcal{P}$  to the boundary of  $\Gamma \backslash D$ . In classical terms the period matrices are polynomials in  $\log t$  with analytic coefficients, and we let  $t \rightarrow 0$ . Lie theory provides a classification of how this may happen [KR].

The other type of subvarieties of  $\overline{\mathcal{P}}$  is that corresponding by the Mumford-Tate sub-domains  $D' \subset D$ . Associated to a polarized Hodge structure  $(V, Q, F)$  is the algebra  $T(F)$  of Hodge tensors in the tensor algebra of  $V$ , and  $D'$  is the orbit in  $D$  of  $T(F)$  under the Lie group associated to the subgroup  $G' \subset G$  preserving that algebra. Geometrically, for algebraic surfaces in first approximation one thinks of those  $X$ 's having additional Hodge classes in  $H^2(X)$ .

Our objective is use the structure of  $\overline{\mathcal{P}}$  to help understand and organize the structure of  $\overline{\mathcal{M}}$ . Perhaps the simplest illustration of this is given by the following

**Model Example:** For algebraic curves the structure of  $\overline{\mathcal{M}}_g$  is a much studied and very beautiful subject. For the first case  $g = 2$  the picture of the stratification is



The results we shall discuss about algebraic surfaces are of the following two types.

1. General results valid for any KSBA moduli space of general type surfaces.
2. Results about  $I$  surfaces, defined to be smooth surfaces  $X$  with  $q(X) = 0$ ,  $p_g(X) = 2$  and  $K_X$  ample. Informally stated we shall see there are three results about the completed moduli space  $\overline{\mathcal{M}}_I$ :
  - (a) for the part  $\overline{\mathcal{M}}_I^G$  of Gorenstein degenerations there is an analogous picture to the solid line part of the one above for  $g = 2$  curves; the stratification of  $\overline{\mathcal{M}}^G$  is faithfully captured by the extended period mapping

$$\overline{\Phi} : \overline{\mathcal{M}}^G \rightarrow \overline{\mathcal{P}};$$

- (b) in a phenomenon not present in the curve case, Hodge theory provides a guide as to how to desingularize a general point of the boundary  $\overline{\mathcal{M}}_g \setminus \mathcal{M}_g$ ;
- (c) for the part  $\overline{\mathcal{M}}_g^{NG}$  of normal, non-Gorenstein degenerations these correspond to the dotted lines in the above figure and there are partial results, an interesting example, and a question/conjecture about what the general story might be.

## Summarizing:

- ▶ For curves,  $\overline{\mathcal{M}} = \overline{\mathcal{M}}_g$  is much studied and much is known about its rich structure.
- ▶ For  $\dim X \geq 2$ , so far as I am aware only a few examples have been partially worked out (cf. [FPR], [H]). In this lecture we will use Hodge theory as a guide to help give some answers to the above question for what is in some sense the “first” general type surface one comes to (an analogue of  $g = 2$  curves).
- ▶ An invariant of  $\mathcal{M}$  is given by the period mapping

$$\Phi : \mathcal{M} \rightarrow \mathcal{P} \subset \Gamma \backslash D.$$

Here, as discussed in Colleen Robles' talk,  $D$  is the period domain parametrizing polarized Hodge structures  $(V, Q, F)$  of a given weight and type,  $\Gamma$  is a discrete subgroup of  $\text{Aut}(V_{\mathbb{Z}}, G)$  that contains the monodromy group (discussed further below).<sup>§</sup>

- ▶ There is a Hodge-theoretically constructed canonical completion  $\overline{\mathcal{P}}$  of  $\mathcal{P}$  and an extension

$$\overline{\Phi} : \overline{\mathcal{M}} \rightarrow \overline{\mathcal{P}}$$

of the period mapping. Our general objective is to use the known structure of  $\overline{\mathcal{P}}$  together with general algebro-geometric methods to infer properties of  $\overline{\mathcal{M}}$ .

---

<sup>§</sup>It has recently been proved in [BBT] that when  $\Gamma$  is arithmetic  $\mathcal{P}$  is an algebraic variety over which the augmented Hodge line bundle  $\Lambda \rightarrow \mathcal{P}$  is ample.

## II. Background and general results

### A. Moduli theory (informal account)

- ▶ We will consider moduli spaces  $\mathcal{M}$  whose points correspond to (equivalence classes of) varieties  $X$  that have the property
  - ▶  $X$  is smooth or has canonical singularities;
  - ▶  $K_X$  is ample.

The first condition means that the Weil canonical divisor class  $K_X$  is a line bundle and that for a minimal desingularization  $\tilde{X} \rightarrow X$  we have  $f^*K_X = K_{\tilde{X}}$ .

- ▶ For this talk there are two main points:
  - (i) the canonical completion  $\overline{\mathcal{M}}$  exists (we will recall its definition);
  - (ii) in the case when  $\dim X = 1, 2$  there is a classification of the singularities of the curves and surfaces corresponding to points of  $\partial\mathcal{M} = \overline{\mathcal{M}} \setminus \mathcal{M}$ .

For (i) we use the valuative criterion: Given a family

$$\mathcal{X}^* \xrightarrow{\pi} \Delta^*$$

of smooth general type varieties  $X_t = \pi^{-1}(t)$ , possibly after a base change we want to define a unique limit  $\lim_{t \rightarrow 0} X_t = X_0$ . The conditions are

- (a)  $mK_{X_0}$  should be a line bundle for some  $m$ ;
- (b)  $X_0$  has semi-log-canonical (slc) singularities;
- (c)  $K_{X_0}$  is ample.

Condition (b) is local along  $X_0$ ; (c) is global.

To explain (b) we consider a minimal completion

$$\begin{array}{ccc} \mathcal{X}^* & \subset & \mathcal{X} \\ \downarrow & & \downarrow \pi \\ \Delta^* & \subset & \Delta. \end{array}$$

Then in 1<sup>st</sup> approximation (b) means that

$\mathcal{X}$  should have canonical singularities.

Varieties  $X_0$  with these properties are said to be *stable* there are a number of highly non-trivial technical issues required to properly formulate much less prove this result. For a discussion of these we refer to [K], [H] and the references cited there.

- ▶ The second point is that for curves and surfaces the singularities of a stable  $X_0$  have been classified.

For simplicity of notation we shall simply use  $X$  instead of  $X_0$ . For curves, the singularities of  $X$  consist of nodes. For surfaces it will be convenient to use a rough organization of the singularity type given by the table

$X$	normal singularities	non-normal singularities
$K_X$	G	NG

where G stands for Gorenstein and NG stands for non-Gorenstein.

In the  $K_X$ -NG spot, by definition there is smallest integer, the *index*  $m \geq 2$  of  $X$ , such that  $mK_X$  is a line bundle.<sup>¶</sup> The entries in the first row mean that the singularities of  $X$  could be isolated (i.e., points), or could occur along curves. In the  $K_X$ -G spot,  $K_X = \omega_X$  is the dualizing sheaf and is a line bundle.

In this talk we shall be particularly interested in the case when  $X$  has normal singularities; we shall denote by  $(X, p)$  the pair given by a stable surface  $X$  and a normal (and hence isolated) singular point  $p$ . Then the classification breaks into 2-types.

---

<sup>¶</sup>A significant issue is to give a good bound on the index. Here we refer to [RU] for interesting recent work.

**K<sub>X</sub>-G:** These include the *canonical singularities*, concerning which there is a rich and vast literature (e.g., Chapter 4 in [R]). They are also referred to as Du Val or ADE singularities and are locally analytically equivalent to isolated hypersurface singularities  $f(x_1, x_2, x_3) = 0$  in  $\mathcal{U} \subset \mathbb{C}^3$ . For example,  $A_n$  is given by

$$x_1^2 + x_2^2 + x_3^{n+1} = 0.$$

For  $n = 1$  there is the standard resolution  $(\tilde{X}, \tilde{C}) \rightarrow (X, p)$  where  $\tilde{C} \cong \mathbb{P}^1$  is a  $-2$  curve (i.e.,  $\tilde{C}^2 = -2$ ). In general the  $\tilde{C}$  is a configuration of  $-2$  rational curves corresponding to the nodes in a Dynkin diagram.

The remaining singularities are the simple elliptic singularities and the cusps. An important non-trivial constraint in the moduli theory of surfaces considered in this talk is that the isolated singularities (normal case) should be smoothable. For simple elliptic singularities this means that the degree  $d = -\tilde{C}^2$  should satisfy  $1 \leq d \leq 9$ .

For the next type we shall use the singularity theorists' notation

$$\frac{1}{n}(1, r), \quad \gcd(n, r) = 1$$

for the quotient  $\mathbb{C}^2 / \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^r \end{pmatrix}$  where  $\zeta = e^{2\pi i/n}$  is a primitive  $n^{\text{th}}$  root of unity.

**K<sub>X</sub>-NG:** These are required to be  $\mathbb{Q}$ -Gorenstein smoothable, meaning that there should be a local smoothing whose relative dualizing sheaf is  $\mathbb{Q}$ -Cartier (cf. [H]). Then there are two types of such singularities:

- (i) the  $\frac{1}{dn^2}(1, dna - 1)$  singularities; for  $d = 1$  these are called *Wahl singularities*. Again for these there is an extensive literature (cf. [H] and the references cited therein);
- (ii) the  $\mathbb{Z}_2$ -quotients of simple elliptic or cusp singularities (cf. (3.24)(c) in [K]).

The non-isolated KSBA singularities are given by pairs  $(X, C)$  where  $C$  is a (possibly reducible) double curve having isolated pinch points and nodes. Typically there is a resolution


$$(\tilde{X}, \tilde{C}) \rightarrow (X, C)$$

where  $\tilde{X}$  is smooth,  $\tilde{C} \subset \tilde{X}$  is a possibly reducible nodal curve with an involution

$$\tau : \tilde{C} \rightarrow \tilde{C},$$


and  $(X, C)$  is the quotient of  $(\tilde{X}, \tilde{C})$  by the involution  $\tau$  where we identify  $p \in \tilde{C}$  with  $\tau(p) \in \tilde{C}$ .

A particularly interesting example of this is due to Liu-Rollenske [LR]. Here  $\tilde{X}$  will be a blow up of  $\mathbb{P}^2$ , and the initial picture is



where we identify  $L_1$  and  $L_2$  by  $\left\{ \begin{array}{l} 12 \longleftrightarrow 21 \\ 13 \longleftrightarrow 24 \\ 14 \longleftrightarrow 23 \end{array} \right\}$  and similarly for  $L_3$  and  $L_4$ .

To define  $\tau$  we must blow up the intersection points  $L_i \cap L_j$



The choice of  $\tau$  is drawn in. Dotted lines are exceptional divisors  $E_{ij}$ . Here  $\tilde{C}$  is a reducible nodal curve. The surface  $X$  is an example of Kollár's gluing construction.

- ▶ Finally we shall be particularly interested in two subvarieties of  $\overline{\mathcal{M}}$ :
  - ▶  $\mathcal{M}_f \subset \overline{\mathcal{M}}$ , defined to be the points corresponding to surfaces that have a smoothing with finite monodromy;
  - ▶  $\overline{\mathcal{M}}^G \subset \overline{\mathcal{M}}$ , defined to be the points corresponding to surfaces  $X$  whose singularities are Gorenstein. In this case  $K_X = \omega_X$  is a line bundle, and duality and Riemann-Roch hold as if  $X$  were smooth.

## B. Hodge theory

- ▶ Roughly speaking Hodge theory has at least the following four interrelated aspects:
- (1) *Topological*: the deeper topological properties of complex algebraic varieties  $X$  arise from the functorial Hodge structure, or mixed Hodge structure, on the cohomology  $H^*(X)$ .
  - (2) *Analytical*: associated to a family  $H_t^n$  of Hodge structures on the punctured disc  $\Delta^*$  there is an essentially unique limiting mixed Hodge structure  $H_{\text{lim}}^n$ ; the topological aspect of the family is reflected by the monodromy

$$T = T_s T_u \quad (\text{Jordan decomposition})$$

where the semi-simple part  $T_s$  is of finite order (the eigenvalues are roots of unity), and the unipotent part  $T_u = e^N$  where  $N^{n+1} = 1$ .

Lie theory and complex analysis combine to give the subtle analytic properties of  $H_{\text{lim}}^m$  (cf. [S] for the 1-parameter case and [CKS] for the several parameter case over  $(\Delta^*)^k$ ).

- (3) *Geometric*: associated to a Hodge structure or to a 1<sup>st</sup> order variation of such there may be algebro-geometric objects; we shall illustrate this with some examples.<sup>||</sup>

**Classical example:** Associated to a principally polarized Hodge structure of weight 1 there is a divisor  $\Theta$ . For the Jacobian of a smooth algebraic curve  $C$  this divisor reflects the geometry of the curve (Riemann), and in fact determines it (Torelli).

<sup>||</sup>These types of applications of Hodge theory to questions in algebraic geometry have, at least recently, been less prevalent than the first two. However, when one is dealing with moduli parameters naturally appear in the geometric picture so one may reasonably anticipate more of the third type of applications.

**Extended classical example:** In the non-classical case when the period domain  $D$  is not Hermitian symmetric there is no analogue of  $\Theta$ . However if we consider the 1<sup>st</sup> order variation of a smooth curve  $C$ , then the 1<sup>st</sup> order Hodge theoretic data gives the space  $I_2(C)$  of quadrics through the canonical curve  $\varphi_K(C) \subset \mathbb{P}^{g-1}$ . For  $C$  non-hyperelliptic  $I_2(C)$  determines  $C$ . This method extends to the non-classical case and leads to various Torelli-type theorems, results about algebraic cycles, etc. (cf. [Gr] and [CM-SP]).

**Non-classical example:** There is quite interesting geometry associated to the extension data in a limiting mixed Hodge structure and its first variation, both in the smoothing and equisingular directions.\*\* This will be illustrated below and discussed in more detail in a later talk.

- (4) *Non-abelian Hodge theory*: this is the study initiated by Simpson of the fundamental groups of algebraic varieties via their linear representation, especially those that arise from variations of Hodge structures. We shall not be able to discuss this very interesting area in this talk.

---

\*\*There is also geometry associated to the extension data in a mixed Hodge structure that arises from a geometric situation (cf. [C]). There is very interesting additional geometry in the case of a *limiting* mixed Hodge structure.

**Stratification of  $\overline{\mathcal{P}}$ :** As mentioned above, there are two basic types of subvarieties of  $\overline{\mathcal{P}}$  and the resulting amalgam of these. The first type is the stratification associated to the boundary components given by the types of limiting mixed Hodge structures that occur when the polarized Hodge structures degenerate. Lie theory provides a classification of how this may happen (cf. [KR]).  
A simple type of Mumford-Tate sub-domains occurs when the PHS decomposes non-trivially into a direct sum of PHS's; these correspond to projection operators in  $\text{End}(V) \subset T(F)$ .

**Generalized stratification of  $\overline{\mathcal{P}}$ :** By generalized stratification we shall mean a set of subvarieties, not necessarily disjoint but whose union is all of  $\overline{\mathcal{P}}$ , and which satisfy certain conditions that will not be spelled out here. The generalized strata will be of the two types discussed above. The first type will be referred to as boundary components; these refer to the LMHS's that appear in  $\overline{\mathcal{P}}$ . The second will be called Mumford-Tate loci.<sup>††</sup>

---

<sup>††</sup>They are also sometimes referred to as a Noether-Lefschetz loci.

**Boundary components:** Very roughly speaking there are two types of boundary components; viz. over  $\mathbb{Q}$  and over  $\mathbb{Z}$ . There is yet to be a formal definition of the latter, which in this talk will be taken to be the  $\mathbb{Q}$ -boundary component together with the  $G_{\mathbb{Z}}$  conjugacy class of  $T_s$  (which is closely related to the *spectrum* in the case of isolated hypersurface singularities). For the purposes of the talk, for the former we use the conjugacy class of  $N$ .

For  $n = 1$  since  $N^2 = 0$  this is determined by rank  $N$ .


For  $n = 2$  one has the classification

- ▶  $N^2 = 0$ ; then we have rank  $N$ .
- ▶  $N^2 \neq 0$ ; then we have rank  $N$  and rank  $N^2$ .

One may picture the  $\mathbb{Q}$ -boundary structure by a diagram in which the conjugacy classes and possible degenerations are represented. For  $n = 1$  and  $h^{1,0} = g$  the diagram is

$$I_0 \text{ --- } I_1 \text{ --- } \dots \text{ --- } I_g.$$

For  $n = 2$  and  $h^{2,0} = 2$  the diagram is



This diagram will be refined when we discuss  $l$ -surfaces. References to these diagrams are given in Colleen's lecture and in [KR].

## C. Some general results:

- ▶ Let  $\mathcal{M}$  be a KSBA moduli space for a class of surfaces of general type and with canonical completion  $\overline{\mathcal{M}}$ . The first general result concerns the period mappings of  $\mathcal{M}$  and  $\overline{\mathcal{M}}$ . It is known (Vakil) that the structure of  $\overline{\mathcal{M}}$  may be arbitrarily nasty and the exact technical conditions under which the following results will hold have not been worked out. We do assume that each component of  $\overline{\mathcal{M}}$  is generically reduced and that a general point corresponds to a smooth surface. Then there is a holomorphic period mapping

$$\Phi : \mathcal{M} \rightarrow \mathcal{P} \subset \Gamma \backslash D$$

whose image  $\mathcal{P}$  is a locally closed analytic subvariety.

From [BBT] it follows that when  $\Gamma$  is arithmetic the closure of  $\mathcal{P}$  in  $\Gamma \backslash D$  is a quasi-projective algebraic variety over which the Hodge line bundle is ample.<sup>‡‡</sup>

- ▶ The first “result” is that the above period mapping extends to

$$(1.1) \quad \bar{\Phi} : \bar{\mathcal{M}} \rightarrow \bar{\mathcal{P}}$$

and that over  $\bar{\mathcal{P}}$  the extended Hodge line bundle is ample. This “result” has been established only in special cases. What is known [GGLR] is that  $\bar{\mathcal{P}}$  exists as a compact Hausdorff space with a stratification by complex analytic subvarieties and that  $\bar{\Phi}$  is defined and is a continuous proper mapping.

---

<sup>‡‡</sup>The interesting work [BBT] uses  $o$ -minimal structures (arising initially from model theory) to put an algebraic structure on  $\mathcal{P}$ . The techniques introduced there and in the references to that work seem certain to have further applications to Hodge theory.

The structure sheaf  $\mathcal{O}_{\overline{\mathcal{P}}}$  is defined to be the sheaf of continuous functions that restrict to be holomorphic on strata. What remains to be proved is that  $\mathcal{O}_{\overline{\mathcal{P}}}$  has enough local functions. This is a global problem along the compact fibres of  $\overline{\Phi}$ .

As a set  $\overline{\mathcal{P}}$  consists of the associated graded PHS's to the equivalence classes of LMHS's obtained from families  $\mathcal{X}^* \rightarrow \Delta^*$  of smooth surfaces parametrized by discs  $g : \Delta^* \rightarrow \mathcal{M}$ . The essential geometric content of the above assertion about  $\overline{\Phi}$  is that

$$\mathrm{Gr} \left( \lim_{t \rightarrow 0} H^2(X_t) \right)$$

depends only on the limit surface  $X_0$  and not on the  $\overline{g} : \Delta \rightarrow \overline{\mathcal{M}}$  extending  $g$  above with  $\overline{g}(0)$  corresponding to  $X_0$ . That is, the associated graded to the LMHS does not depend on the particular smoothing of  $X_0$  (there may be several components of such).

In the example of  $l$ -surfaces discussed one may see this directly.

- ▶ For the next general result we recall the notation

$$\mathcal{M}_f \subset \overline{\mathcal{M}}$$

for the subvariety of  $\overline{\mathcal{M}}$  parametrizing singular surfaces  $X$  such that there exists a smoothing  $\mathcal{X} \rightarrow \Delta$  of  $X = X_0$  with finite monodromy. Then

*The period mapping extends to  $\Phi : \mathcal{M}_f \rightarrow \Gamma \backslash D$ .*

Moreover,

$$\overline{\mathcal{M}}^{\text{NG}} \subset \mathcal{M}_f.$$

Here,  $\overline{\mathcal{M}}^{\text{NG}}$  denotes the subvariety of  $\overline{\mathcal{M}}$  parametrizing *normal* surfaces  $X$  having non-Gorenstein singularities. Informally stated, to a normal and smoothable surface  $X$  having non-Gorenstein semi-log-canonical singularities one may associate a pure polarized Hodge structure  $H_{\text{lim}}^2(X)$ .

This latter result is more of an observation than a theorem: it is a consequence of the statements

- ▶ normal surfaces with rational singularities are parametrized by a subvariety of  $\mathcal{M}_f$  (i.e., they have finite monodromy);\*

---

\*From a Hodge theoretic perspective this result is basically obvious. The limit  $\tilde{\omega}$  of holomorphic 2-forms  $\omega_t \in H^0(\Omega_{X_t}^2)$  will have  $\tilde{\omega} \in H^0(\Omega_{\tilde{X}}^2(\tilde{C}))$  where  $(\tilde{X}, \tilde{C})$  is a desingularization of  $X$ . Then  $\text{Res } \tilde{\omega}$  will be a 1-form on a set of  $\mathbb{P}^1$ 's with opposite residues at intersection points. Since a component of  $\tilde{C}$  will be a tree, not a cycle, of  $\mathbb{P}^1$ 's, we see that  $\tilde{\omega} \in H^0(\Omega_{\tilde{X}}^2)$ . The result follows from the Clemens-Schmid exact sequence ([M]).

and from the above classification of normal, slc singularities

- ▶ normal, non-Gorenstein slc singularities are rational.

Canonical singularities are also rational so they are also parametrized by a subvariety in  $\mathcal{M}_f$ . For  $I$ -surfaces thus far there is no example other than the ones mentioned above of singular surfaces with finite monodromy. It follows from the classification in [FPR] that if  $X$  is Gorenstein and non-normal, then the monodromy is infinite (i.e.,  $N \neq 0$ ). A natural question is whether non-normal  $I$ -surfaces always have infinite monodromy?

- ▶ The normal surfaces  $X$  with infinite monodromy are those with either simple elliptic singularities or cusps. For the following result we assume that a general smooth surface is regular and that the singular surface has either  $e$  simple elliptic singularities or  $c$  cusps but does not have some of each.<sup>†</sup> The statement is

$$\begin{cases} \text{rank } N \leq e \leq \rho_g + 1 \\ \text{rank } N^2 \leq c \leq \rho_g + 1. \end{cases}$$

Below we will give a Hodge-theoretic proof of the first part of this result.

- ▶ Finally we will explain a general statement that might hold and that can be established in a couple of cases. The period mapping extends from  $\mathcal{M}$  to give

$$\Phi : \mathcal{M}_f \rightarrow \Gamma \backslash D.$$

---

<sup>†</sup>There is a general result without these assumptions, but it is more involved to formulate and the special cases given here capture the essential geometric content of the general result.

As noted earlier that the image  $\mathcal{P}$  is a *closed* analytic subvariety and it follows from the results in [BBT] that  $\mathcal{P}$  is quasi-projective.

Let  $M \subset \mathcal{M}_f$  be an irreducible component of  $\mathcal{M}_f$ . The question that arises is

*Does there exist a  $\Gamma$ -invariant Mumford-Tate subdomain  $D' \subset D$  with  $\Gamma'$  the discrete group of automorphisms of  $D'$  induced by  $\Gamma$  such that*

$$M = \Phi^{-1}(\mathcal{P} \cap (\Gamma' \backslash D'))?$$

Informally this means that these components of moduli can be detected Hodge theoretically.

How might one prove this, at least in some special cases? For those  $M$  such that the surfaces  $X$  parametrized by  $M$  are normal with either canonical or non-Gorenstein singularities, such singularities are rational and the resolution

$$(\tilde{X}, \tilde{C}) \rightarrow (X, p)$$

of a particular one has for  $\tilde{C}$  a configuration of  $\mathbb{P}^1$ 's.

Recalling that  $X$  gives a PHS  $\Phi(X) \in \Gamma \setminus D$ , the  $\mathbb{P}^1$ 's give Hodge classes that may not be present on a general point in  $D$ , and then  $D'$  could be the Mumford-Tate domain defined by PHS's having these additional Hodge classes.

One then hopes to use a variational argument to show that in  $T \text{Def } X$  the condition to retain these Hodge classes defines the tangent space to  $M \subset \overline{\mathcal{M}}$ . This argument has been carried out in the two cases

- ▶ an  $A_1$ -singularity that is not a base point of  $|K_X|$ ;
- ▶ the  $\frac{1}{4}(1, 1)$ -singularity on the general  $I$ -surface having that type of singularity.

In both use is made of differential of the period mapping at a singular surface that will be discussed elsewhere.

- ▶ To illustrate how Hodge theory and algebraic geometry interact and complement one another we shall give a proof of the result stated above:

*Let  $X$  be a normal surface having  $e$  simple elliptic singularities  $p_i$  and that is a KSBA degeneration of smooth regular general type surface with geometric genus  $p_g$ . Then*

$$\text{rank } N \leq e \leq p_g + 1.$$

*Moreover, if equality holds on the right, then  $\text{rank } N = p_g$  and all of the elliptic curves  $\tilde{C}_i$  that contract to  $p_i$  under the minimal desingularization of  $X$  are isogeneous to a fixed elliptic curve.*

**Proof:** We set  $\tilde{C} = \sum \tilde{C}_i$  and denote by  $(\tilde{X}, \tilde{C})$  the minimal desingularization of  $(X, \{p_i\})$ . The cohomology sequence of the Poincaré residue sequence  $0 \rightarrow \Omega_{\tilde{X}}^2 \rightarrow \Omega_{\tilde{X}}^2(\tilde{C}) \rightarrow \omega_{\tilde{C}} \rightarrow 0$  gives

$$0 \rightarrow H^0(\Omega_{\tilde{X}}^2) \rightarrow H^0(\Omega_{\tilde{X}}^2(\tilde{C}_1 + \cdots + \tilde{C}_k)) \rightarrow \bigoplus_{i=1}^k H^0(\Omega_{\tilde{C}_i}^1) \rightarrow H^1(\Omega_{\tilde{X}}^2) \rightarrow 0$$

$$p_g(\tilde{X})$$

$$h^0(K_X) = p_g$$

$$k$$

$$q(X)$$

where the dimensions appear below the groups in question. The zero on the right is due to  $h^2(K_X) = 0$  since  $X$  is a KSBA degeneration of a regular surface. This gives

$$k = p_g + q(\tilde{X}) - p_g(\tilde{X}).$$

By Castelnuovo's lemma [BPVdV], for  $\alpha$  and  $\beta \in H^0(\Omega_{\tilde{X}}^1)$

$\alpha \wedge \beta = 0 \implies \alpha$  and  $\beta$  pull back from a curve of genus  $\geq 2$ .

If the 1-forms on  $\tilde{X}$  do not pull back from such a curve  $C$ , then for  $\alpha_1, \dots, \alpha_q$  a basis for  $H^0(\Omega_{\tilde{X}}^1)$  the 2-forms  $\alpha_1 \wedge \alpha_2, \dots, \alpha_1 \wedge \alpha_q$  are linearly independent and

$$p_g(\tilde{X}) \geq q(\tilde{X}) - 1$$

which gives

$$(2.1) \quad k \leq p_g + q(\tilde{X}) - (q(\tilde{X}) - 1) \leq p_g + 1.$$

If on the other hand there is  $f : \tilde{X} \rightarrow C$ , then the dual to  $H^0(\Omega_C^1) \xrightarrow{f^*} H^0(\Omega_{\tilde{X}}^1)$  is a map

$$H^2(\Omega_{\tilde{X}}^1) \rightarrow H^1(\mathcal{O}_C).$$

The Gysin map

$$\bigoplus_{i=1}^k H^0(\Omega_{\tilde{C}_i}^1) \rightarrow H^2(\Omega_{\tilde{X}}^1)$$

is dual to the restriction map  $H^0(\Omega_{\tilde{X}}^1) \rightarrow \bigoplus_{i=1}^k H^0(\Omega_{\tilde{C}_i}^1)$ . Then using the  $f^*$  above

$$(2.2) \quad H^0(\Omega_C^1) \rightarrow H^0(\Omega_{\tilde{C}_i}^1)$$

is zero unless  $g(C) = 1$  and  $\tilde{C}_i \xrightarrow{f} C$  is an isogeny. Thus if  $g(C) \geq 2$ , the mappings (2.2) are all zero. As a consequence

$\alpha \in H^0(\Omega_{\tilde{X}}^1)$  and  $\alpha|_{\tilde{C}_i} \neq 0 \implies \alpha \wedge \beta \neq 0$  for  $\beta \in H^0(\Omega_{\tilde{X}}^1)/\mathbb{C}\alpha$ .

It follows that for

$$m = \dim \left( \text{Im} \{ H^0(\Omega_{\tilde{X}}^1) \rightarrow \bigoplus_{i=1}^k H^0(\Omega_{C_i}^1) \} \right)$$

we have

$$\rho_g(\tilde{X}) \geq m - 1,$$

which then by (2.1) gives  $k \leq \rho_g + 1$ .

The lower bound

$$\text{rank } N \leq k$$

follows from the Clemens-Schmid exact sequence (cf. [M]).

**Remark:** This argument may be adapted to show that if  $X$  has a cusp, then

$$\text{rank } N^2 \leq c \leq \rho_g + 1.$$

Each cusp is a cycle of some number  $\ell$  of  $\mathbb{P}^1$ 's, and we do not know how to bound the  $\ell$ .

### III. $I$ -surfaces

- ▶ Recall that an  $I$ -surface is defined to be a smooth (or having canonical singularities) surface  $X$  that satisfies
  - ▶  $X$  is minimal of general type
  - ▶  $q(X) = 0$  and  $p_g(X) = 2$
  - ▶  $K_X^2 = 1$

It appears that regular surfaces for which Noether inequality

$$p_g(X) \leq \frac{K_X^2}{2} + 2$$

is close to equality seem to have favorable qualities for the use of Hodge theory to study their moduli, and some of what follows has also been carried out for  $H$ -surfaces satisfying the first two conditions above together with  $K_X^2 = 2$  ( $H$  stands for Horikawa [Ho] who made an extensive analysis of surfaces with small  $c_1^2$ ).

Informally stated parts of the results that we shall discuss are

- ▶  $\overline{\Phi}, \overline{\mathcal{M}}_I^G \rightarrow \overline{\mathcal{P}}$  is a mapping of stratified varieties that is bijective on components;
- ▶ the extension data in the limiting mixed Hodge structures over  $\overline{\mathcal{P}} \setminus \mathcal{P}$  provides a guide for desingularizing  $\overline{\mathcal{M}}_I^G$  over the boundary of  $\mathcal{M}_I$ .

As mentioned above, an extensive analysis of  $\overline{\mathcal{M}}_I^G$  has been carried out by [FPR]; this will be summarized in a table below and discussed in more detail in another lecture. Here we shall first give a Lie-theoretically constructed table of the possible types of LMHS's that could appear in  $\overline{\mathcal{P}} \setminus \mathcal{P}$ . We will then give the FPR table in the normal Gorenstein case and sketch how the Hodge-theoretic table suggests and interprets the algebro-geometric one.

In a subsequent talk we hope to present more detailed discussion of how the Hodge theoretic and algebro-geometric approaches may be combined and interpreted for both normal and non-normal singularities and in both the  $I$ -surface and  $H$ -surface cases.

We emphasize that this is work in progress and some of the details remain to be worked through. The main missing piece is the analysis of the non-Gorenstein case; specifically, for  $I$ -surfaces are there examples beyond the  $\frac{1}{4}(1, 1)$  Wahl singularity (and thus is the index at most equal to 2)?

- ▶ The principal properties of  $I$ -surfaces we shall use are
  - ▶  $h^1(mK_X) = 0$  for  $m \geq 0$  and

$$(3.1) \quad h^0(mK_X) = \begin{cases} 2 & \text{for } m = 1 \\ \frac{m(m-1)}{2} + 2, & m \geq 2; \end{cases}$$

- ▶ using Kawamata-Vieweg vanishing one sees that these properties hold for any Gorenstein  $I$ -surface where the Weil canonical divisor class and the dualizing sheaf  $\omega_X$  coincide as line bundles;
- ▶ in the Gorenstein case the pluri-canonical ring

$$R(X) = \bigoplus^m H^0(mK_X)$$

has the *postulated form*, meaning that generators and relations are added only when required by (3.1) (cf. [FPR] for a proof);

- ▶ classically from Castelnuovo-Enriques and since the work of Bonbieri and others one studies general type surfaces via their pluri-canonical maps ([BPVdV])

$$\varphi_{mK_X} : X \rightarrow \mathbb{P}H^0(mK_X)^* \cong \mathbb{P}^{P_m-1};$$

- ▶ instead of these it is frequently more convenient to use weighted projective spaces corresponding to when new generators are added; thus

$$\varphi_{K_X} : X \dashrightarrow \mathbb{P}^1 \quad (|K_X| = \text{pencil of hyperelliptic curves})$$

$$\varphi_{2K_X} : X \rightarrow \mathbb{P}(1, 1, 2) \rightarrow \mathbb{P}^3 \quad (|2K_X| \text{ is base-point-free})$$


⋮

$$\varphi_{5K_X} : X \hookrightarrow \mathbb{P}(1, 1, 2, 5) \hookrightarrow \mathbb{P}^{12} \quad (|5K_X| \text{ is very ample}).$$

We denote by  $(x_0, x_1, y)$  weighted homogeneous coordinates in  $\mathbb{P}(1, 1, 2)$ , and by  $(x_0, x_1, y, z)$  those for  $\mathbb{P}(1, 1, 2, 5)$ .

▶ Equations/picture

▶  $z^2 = F_5(t_0, t_1, y)z + F_{10}(t_0, t_1, y)$

▶   $\mathbb{P}(1, 1, 2) \hookrightarrow \mathbb{P}^3$  given by  
 $[x_0, x_1, y] \rightarrow [x_0^2, x_0x_1, x_1^2, y]$

$X = 2:1$  map branched over  $P$  and  $V \in |\mathcal{O}_{\mathbb{P}^3}(5)|$   
 where  $V$  does not pass through the vertex  $P^\ddagger$

▶ The moduli of smooth  $X$ 's may be analyzed

- ▶ from the above equation;
- ▶ cohomologically using the Jacobian ideal formalism (cf. [Gr]) for weighted projective spaces.

---

$^\ddagger$ Any Gorenstein  $X$  is irreducible ( $K_X^2 = 1$ ) and is given by such a picture where  $V$  does not contain  $P$ .

Each has its advantages. The first is useful in studying degenerations, and also possibly in using GIT where in this case the group is non-reductive. From the second approach one has

- ▶  $\mathcal{M}$  is reduced and smooth of dimension 28;
- ▶ local Torelli holds.

More precisely, for smooth  $X$  the differential of the period mapping is 1-1. Versions of this have been proved by Carlson-Toledo [CT] and Pearlstein-Zhang [(PZ)].

- ▶ We recall that for any non-classical period domain  $D$  there is a non-trivial invariant distribution  $I \subset TD$  (the infinitesimal period relation or IPR) such that any period mapping

$$\Phi : B \rightarrow \Gamma \backslash D$$

satisfies

$$\Phi_* : TB \rightarrow I.$$

For polarized Hodge structures of weight 2 and with  $h^{2,0} = 2$

$\dim D = 2h^{1,1} + 1$  and  $l$  is a contact distribution.

From Noether's formula

$$\chi(\mathcal{O}_X) = \frac{1}{12}(c_1^2 + c_2)$$

we may infer that  $h^{1,1} = \dim H^{1,1}(X)_{\text{prim}} = 28$ .

From this it follows that

$\Phi(\mathcal{M}) \subset \Gamma \setminus D$  is a contact submanifold.

▶ Remark that a cohomological analysis also gives

▶  $h^0(T_X) = 0$  (since  $X$  is general type)

▶  $\chi(T_X) = 28$  (Hirzebruch-Riemann-Roch)

⇒  $h^2(T_X) = 0$ ,

which again shows that for  $X$  smooth the Kuranishi space is smooth of dimension 28. We suspect that this still holds when  $X$  has canonical singularities, but this has not been checked.

▶ Finally the classical methods of Lefschetz ([L] and [B]) may be adapted to show that

the monodromy group  $\Gamma = \Phi_* (\pi_1(\mathcal{M}_{l,\text{reg}}))$  is arithmetic.

It is not known if  $\Gamma = G_{\mathbb{Z}}$  is the full arithmetic group. §

---

§Note that since  $K_X^2 = 1$ , the primitive decomposition is

$$H^2(X, \mathbb{Z}) = \mathbb{Z} \cdot c_1(X) \oplus H^2(X, \mathbb{Z})_{\text{prim}}$$

and the intersection form on the primitive part is unimodular.

Also outstanding are the questions

- ▶ Does generic global Torelli hold (i.e., does  $\Phi$  have degree one)?
- ▶ Does global Torelli hold?

We expect that the first question may be addressed using the methods in [Gr]. There are multiple properties (cohomological, geometric using versions of generic global Torelli for boundary components of  $\overline{\mathcal{M}}_g$ ) that we hope to discuss in another talk.

- ▶ Turning to the classification we first give a refined picture of the Hodge-theoretic boundary components of  $\overline{\mathcal{P}}$  for the case of surfaces where  $p_g = 2$ . For this we will use Hodge diamonds to depict the  $\text{Gr}(\text{LMHS})$ . The numbers above some of the entries are dimensions; from those that are depicted all of the remaining dimensions may be determined.



To illustrate how one uses Hodge theory to suggest what to look for in analyzing the stratification of  $\overline{\mathcal{M}}_g^G$ , under a degeneration

$$0 \rightarrow I,$$

a pure Hodge structure of weight 2 with Hodge numbers  $(2, h^{1,1}, 2)$  degenerates into a limiting mixed Hodge structure with graded pieces

$$H_{\text{lim}}^1, H_{\text{lim}}^2, H_{\text{lim}}^1, (-1)$$

where  $H_{\text{lim}}^1$  has Hodge numbers  $(1, 1)$  and  $H_{\text{lim}}^2$  has Hodge numbers  $(1, h^{1,1}, 1)$ . Algebrao-geometrically one may use semi-stable reduction (SSR) to expect that the KSBA degeneration  $X_t \rightarrow X_0$  becomes a family whose central fibre is a normal crossing surface  $X = X_1 \cup \cdots \cup X_m$ , one component of which is the minimal desingularization  $\tilde{X}$  of  $X_0$ .

Since  $N^2 = 0$ , one may also hope that  $X$  has only a smooth double curve  $C$  (no triple points). Moreover, since the  $p_g$  drops by one in the limit, one may reasonably expect that  $\lim \omega_t = \tilde{\omega}$  where  $\omega_t \in H^0(\Omega_{X_t}^2)$  and  $\tilde{\omega} \in H^0(\Omega_{\tilde{X}}^2(\tilde{C}))$  with  $\text{Res}_{\tilde{C}}(\tilde{\omega}) \in H^0(\Omega_{\tilde{C}}^1) = H_{\text{lim}}^{1,0}$ . Thus we may expect that under degenerations of type I the limit surface  $X_0$  has one simple elliptic singularity.

The following is a table of normal Gorenstein degenerations of  $I$ -surfaces with  $N^2 = 0$ . In it

- ▶  $k = \#$  of simple elliptic singularities;
- ▶  $d_i =$  degree of the elliptic singularity;
- ▶ the numbers in the subscripts on I, III are the degrees of the elliptic singularities.

stratum	dimension	minimal resolution $\tilde{X}$	$\sum_{i=1}^k (9 - d_i)$	$k$	$\text{codim}$ in $\overline{\mathcal{M}}_l$
$I_0$	28	canonical singularities	0	0	0
$I_2$	20	blow up of a K3-surface	7	1	8
$I_1$	19	minimal elliptic surface with $\chi(\tilde{X})=2$	8	1	9
$III_{2,2}$	12	rational surface	14	2	16
$III_{1,2}$	11	rational surface	15	2	17
$III_{1,1,R}$	10	rational surface	16	2	18
$III_{1,1,E}$	10	blow up of an Enriques surface	16	2	18
$III_{1,1,2}$	2	ruled surface with $\chi(\tilde{X})=0$	23	3	26
$III_{1,1,1}$	1	ruled surface with $\chi(\tilde{X})=0$	24	3	27

Note that the last column is the sum of the two columns preceding it. This will be illustrated below and more fully explained in another talk. The geometric point will be that the  $\sum_{i=1}^k (9 - d_i)$  term will be the number of parameters in the extension data of the LMHS; this extension data will suggest how to blow up  $\overline{\mathcal{M}}_l$  along the corresponding component to obtain a desingularization of  $\overline{\mathcal{M}}_l$  along that component.

We will also illustrate how Hodge theory and standard algebro-geometric techniques may be combined to identify which K3 surface, elliptic surface, rational and ruled surfaces appear in the table. For example, for  $I_2$  the K3 will have a degree 2 polarization and  $\tilde{C}$  will be the normalization of a tangent to the sextic in  $\mathbb{P}^2$  that is the branch curve for the double covering of the K3. For  $I_1$  the elliptic surface will be an elliptic pencil with  $p_g = 1$  and having a bi-section.

Referring to the above diagram we note that

$$7 = \left\{ \begin{array}{l} \text{number of points to blow up on a} \\ \text{cubic curve } C \text{ in } \mathbb{P}^2 \text{ to obtain a del} \\ \text{Pezzo } \tilde{\mathbb{P}}^2 \text{ containing } \tilde{C} \text{ with } \tilde{C}^2 = 2 \end{array} \right\}$$

$$= 1 + \text{codimension in moduli of } I_2.$$

This suggests that to desingularize  $\overline{\mathcal{M}}_1$  along  $I_2$  we insert normal crossing surfaces  $\tilde{X} \cup_{\tilde{C}} \tilde{\mathbb{P}}^2$ .

These 7 points also give the extension data in the LMHS. Of course this heuristic dimension count must take into account automorphisms etc., and this can be done. The point here is just to suggest how the Hodge theory and geometry interact.

We will also see that using the results in [Ch] and [F2] there are generic global Torelli theorems for the boundary components  $I_2$ ,  $I_1$ , and it may be possible that the methods in [F2] can be adapted to infer a generic global Torelli result for  $I$  surfaces.


We conclude with a brief remark on the non-normal Liu-Rollenske example mentioned earlier. It turns out that there are three irreducible components of  $\overline{\mathcal{M}}_1^G$  whose general member is non-normal and obtained from  $(\tilde{X}, \tilde{C}, \tau)$  by passing to the quotient. For one of these, say  $\mathcal{R}$ , we have

- ▶  $\tilde{X} = \mathbb{P}^2$ ;
- ▶  $\tilde{C} = \text{plane quartic}$ ;
- ▶ for general  $\tilde{C}$  we have  $\tilde{C}/\tau$  is an elliptic curve;
- ▶  $\dim \mathcal{R} = 3$ .

The LMHS's are of types IV and V. Those of type V are Hodge-Tate and the summand on which  $N^2$  is an isomorphism has graded pieces

$$H_{\text{lim}}^0, H_{\text{lim}}^0(-2)$$

where  $h^0 = 2$ . The picture of the boundary component is



There are three parameters in  $\text{Ext}_{\text{MHS}}^1(H_{\text{lim}}^0, H_{\text{lim}}^0(-2))$  and in this case the extension data generically determines the point of  $\mathcal{R}$ . The extreme case is the aforementioned LR surface

$$z^2 = y(x_0^2 - y)^2(x_1^2 - y)^2$$

obtained by identifying opposite sides of a quadrilateral in  $\mathbb{P}^2$ . In this case the LMHS is split.

## References

- [FPR] {
- ▶ M. Franciosi, R. Pardini, and S. Rollenske, *Computing invariants of semi-log-canonical surfaces*, Math. Z. 280 no. 3-4 (2015), 1107–1123.
  - ▶ M. Franciosi, R. Pardini and S. Rollenske, *Gorenstein stable surfaces with  $K_X^2 = 1$  and  $p_g > 0$* , Math. Nachr. 290 (2017), no. 5-6, 794–814.
  - ▶ M. Franciosi, R. Pardini and S. Rollenske, *Log-canonical pairs and Gorenstein stable surfaces with  $K_X^2 = 1$* , Compos. Math. 151 (2015), no. 8, 1529–1542.

- [A] V. I. Arnold, V. V. Goryunov, O. V. Lyashko, V. A. Vasil'ev, *Singular Theory I*, Springer-Verlag, 1988.

[BBT] B. Bakker, Y. Brunebarbe, and J. Tsimerman, *0-minimal GAGA and a conjecture of Griffiths*, 2018, arXiv:1811.12230.

[B] A. Beauville, *Le group de monodromie des familles universelles d'hypersurfaces et d'intersections completes* (French) [The monodromy group of universal families of hypersurfaces and complete intersections], in *Complex Analysis and Algebraic Geometry* (Göttingen, 1985), 8–18, Lect. Notes in Math. **1194**, Springer-Verlag, Berlin, 1986.

[BPVdV] W. Barth, C. Peters, and A. Van den Van, *Compact Complex Surfaces*, Springer-Verlag, New York, 1984.

[C] J. Carlson, *Extensions of mixed Hodge structures* (Journ. Geom. Alg. d'Angers, July 1979, A. Beauville Ed., Sijthoff and Noordhoff, 1980).

- [CM-SP] J. Carlson, S. Müller-Stach, and C. Peters, *Period mappings and period domains*, Cambridge Studies in Advanced Mathematics, vol. 168, Cambridge University Press, Cambridge, 2017, Second edition of [MR2012297]. MR 3727160
- [CKS] E. Cattani, A. Kaplan, and W. Schmid, *Degeneration of Hodge structures*, Ann. of Math. **123** (1986), no. 3, 457–535. MR 0840721
- [Ch] K. Chakris, *The Torelli problem for elliptic pencils*, in *Topics in Transcendental Algebraic Geometri* (P. Griffiths, ed.), 157–182, Princeton Univ. Press, Princeton, NJ, 1984.
- [F1] R. Friedman, *Global smoothings of varieties with normal crossings*, Ann. of Math. (2) **118** (1983), no. 1, 75–114. MR 0707162
- [F2] R. Friedman, *A new proof of the global Torelli theorem for K3 surfaces*, Ann. of Math. (2) **120** (1984), no. 2, 237–269. MR 0763907

[G]

- ▶ P. Griffiths, *Hodge theory and moduli*, Lectures given at the University of Haifa in May 2019.  
<https://hdl.handle.net/20.500.12111/7865>
- ▶ P. Griffiths, *Moduli and Hodge theory*, talk at the University of Illinois at Chicago on April 5, 2019.  
<https://albert.ias.edu/handle/20.500.12111/7868>
- ▶ P. Griffiths, *Using Hodge theory to detect the structure of a compactified moduli space*, talk given on November 27, 2019 at the IMSA conference held at IMATE at UNAM, Mexico City.  
<https://hdl.handle.net/20.500.12111/7873>
- ▶ P. Griffiths, *Hodge theory and moduli*, talk given at the Algebraic Geometry and Applications Conference in St. Petersburg, Russia in May 2018.  
<https://albert.ias.edu/handle/20.500.12111/7864>

- [Gr] M. Green, *Infinitesimal methods in Hodge theory*, in *Algebraic Cycles and Hodge Theory*, Lectures given at the 2nd Session of the Centro Internazionale Matematico Estivo (C.I.M.E) held in Torino, Italy, June 21–29, 1993, (A. Albano, F. Bardelli, eds.), 1–92, Springer-Verlag, Berlin.
- [H] P. Hacking, *Compact spaces of surfaces of general type*, in *Compact Moduli Spaces and Vector Bundles*, Contemp. Math. **564**, 1–18, Amer. Math. Soc., Providence, RI, 2012.
- [Ho] E. Horikawa, *Algebraic surfaces of general type with small  $c_1^2$ . IV*, Invent. Math. **50** (1979), 103–128.
- [KU] K. Kato and S. Usui, *Classifying spaces of degenerating polarized Hodge structures*, Ann. of Math. Stud. **169**, Princeton Univ. Press, Princeton, NJ (2009).

- [KR] M. Kerr and C. Robles, *Variations of Hodge structure and orbits in flag varieties*, Adv. in Math. **315** (2017), 27–87.
- [K] J. Kollár, *Moduli of varieties of general type*, Handbook of moduli. Vol. II, Adv. Lect. Math. (ALM), vol. 25, Int. Press, Somerville, MA, 2013, pp. 131–157. MR 3184176
- [KSB] J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), no. 2, 299–338. MR 0922803
- [L] S. Lefschetz, *L'analyse Situs et la Géométrie Algébrique*, Gauthier-Villars, Paris, 1924.

- [LR] W. Liu and S. Rollenske, *Geography of Gorenstein stable log surfaces*, Trans. Amer. Math. Soc. **368** (2016), no. 4, 2563–2588. MR 3449249
- [M] D. R. Morrison, *The Clemens-Schmid exact sequence and applications*, in *Topics in Transcendental Algebraic Geometri* (P. Griffiths, ed.), 101–120, Princeton Univ. Press, Princeton, NJ, 1984.
- [PS] C. A. M. Peters and J. H. M. Steenbrink, *Mixed Hodge structures*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 52, Springer-Verlag, Berlin, 2008. MR 2393625

- [PZ] G. Pearlstein and Z. Zhang, *A generic global Torelli theorem for certain Horikawa surfaces*, arXiv:1702.06204v1[math.AG] 20 Feb 2017.
- [R] M. Reid, *Chapters on algebraic surfaces*, *Complex Algebraic Geometry*, J. Kollár (ed.), IAS/Park City Math. Series, Vol. 3, Amer. Math. Soc./IAS.
- [RU] J. Rana and G. Urzúa, *Optimal bounds for T-singularities in stable surfaces*, *Adv. Math.* **345** (2019), 814–844. MR 3901675
- [S] W. Schmid, *Variation of Hodge structure: the singularities of the period mapping*, *Invent. Math.* **22** (1973), 211–319. MR 0382272